

Who To Test

Patients

**Fever > 100.4 AND
-New Cough or
-New Shortness of
Breath**

Is the Patient High Risk?

- Hospitalized w/respiratory symptoms
- SNF Residents
- Health Care Workers
- Public Safety Workers
- High Risk* contact w/COVID positive cases
- Homeless patients
- Special Populations:
 - Adults over the age of 60
 - Chronic lung diseases
 - Pregnancy
 - Hemodialysis
 - Immunocompromised such as:
 - Advanced HIV (CD4<200)
 - Cancer therapy
 - Transplant
 - Immunosuppressant drugs

Yes

High Risk Patient Instructions for Home Isolate

- Stay in a room away from other people.
- Use separate bathroom
- Wear a mask when around other people. Household contacts wear mask also when interacting, if possible
- Clean hands after coughing, sneezing, using bathroom, and before eating
- Avoid touching face
- Avoid sharing household items
- Clean high touch surfaces daily
- For more specifics, please reference [CDC Guidelines](#)

Order test

TEST
COVID-19
[Click this Link](#) to
Specimen Collection

Negative COVID-19

**Discontinue Isolation 24
hours after symptom
resolution.**

Positive COVID-19

**Isolate for a minimum of
7 days from the onset of
symptoms
OR
72 hours after symptom
resolution, whichever is
*longer.***

No

ALL OTHERS

Isolate for a minimum of 7
days from the onset of
symptoms
OR
72 hours after symptom
resolution, whichever is
longer.

***High Risk Contact: Within approximately 6 feet
of a COVID-19 case for a prolonged period of
time; close contact can occur while caring for,
living with, visiting, or having direct contact
with infectious secretions of a COVID-19 case.**

Questions? Email:
KPWACOV19Communication@kp.org